

FAMILIES OF SOLUTIONS OF A CUBIC DIOPHANTINE EQUATION

Marc Chamberland

Department of Mathematics and Computer Science, Grinnell College, Grinnell, IA 50112

E-mail: chamberl@math.grin.edu

(Submitted August 1998-Final Revision February 1999)

This study started with an unusual advertisement which appeared (January 6, 1996) in *The Globe and Mail*, Canada's national newspaper. Vivikanand Kadarnauth (of Toronto) presented the "first few cases" in a family of solutions to the "cubic version of the Pythagorean equation"

$$a^3 + b^3 + c^3 = d^3 \tag{1}$$

as

$$\begin{aligned} 4^3 + 5^3 + 3^3 &= 6^3, & 4^3 + 17^3 + 22^3 &= 25^3, \\ 16^3 + 23^3 + 41^3 &= 44^3, & 16^3 + 47^3 + 108^3 &= 111^3, \\ 64^3 + 107^3 + 405^3 &= 408^3, & 64^3 + 155^3 + 664^3 &= 667^3. \end{aligned}$$

Mr. Kadarnauth then asked the reader to find the general pattern. Some of the patterns indicate that the general solution is

$$(a, b, c, d) = (2^{2m}, 2 \cdot 2^{2m} - 3 \cdot 2^m + 3, 2^{3m} - 2 \cdot 2^{2m} + 3 \cdot 2^m - 3, 2^{3m} - 2 \cdot 2^{2m} + 3 \cdot 2^m)$$

and

$$(a, b, c, d) = (2^{2m}, 2 \cdot 2^{2m} + 3 \cdot 2^m + 3, 2^{3m} + 2 \cdot 2^{2m} + 3 \cdot 2^m, 2^{3m} + 2 \cdot 2^{2m} + 3 \cdot 2^m + 3),$$

where m varies over the positive integers. One may generalize this by replacing 2^m with x , thus yielding the one-parameter polynomial families of solutions

$$(a, b, c, d) = (x^2, 2x^2 - 3x + 3, x^3 - 2x^2 + 3x - 3, x^3 - 2x^2 + 3x) \tag{2}$$

and

$$(a, b, c, d) = (x^2, 2x^2 + 3x + 3, x^3 + 2x^2 + 3x, x^3 + 2x^2 + 3x + 3). \tag{3}$$

The second family (3) is equivalent to (2). This is seen by replacing x with $-x$ in (2) and rearranging the terms, since $a^3 + b^3 + c^3 = d^3$ implies $a^3 + b^3 + (-d)^3 = (-c)^3$. By letting $x = v/u$ in (3) and multiplying by u^3 gives the family of solutions listed by Jandasek (see [3], p. 559):

$$(a, b, c, d) = (uv^2, 3u^2v + 2uv^2 + v^3, 3u^3 + 3u^2v + 2uv^2, 3u^3 + 3u^2v + 2uv^2 + v^3). \tag{4}$$

The cubic Diophantine equation (1) has been studied for over 400 years. In 1591, P. Bungus (see [3], p. 550) found the smallest positive solution mentioned above, namely

$$4^3 + 5^3 + 3^3 = 6^3, \tag{5}$$

the same year that Vieta found a family of solutions. (Perelman writes on page 139 in [7]: "It is said that [equation (5)] highly intrigued Plato.") Almost 200 years later, Euler (see [3], p. 552) found that the general **rational** solution to equation (1) may be represented (see [6], p. 292) as

$$\begin{aligned}
 (a, b, c, d) = & (\sigma(-(\xi - 3\eta)(\xi^2 + 3\eta^2) + 1), \\
 & \sigma((\xi^2 + 3\eta^2)^2 - (\xi + 3\eta)), \\
 & \sigma((\xi + 3\eta)(\xi^2 + 3\eta^2) - 1), \\
 & \sigma((\xi^2 + 3\eta^2)^2 - (\xi - 3\eta))),
 \end{aligned} \tag{6}$$

where $\sigma, \xi,$ and η are rationals. The variable σ is simply a scaling factor reflecting the homogeneity of equation (1). Ramanujan [2] also gave a general solution as

$$(a, b, c, d) = (\alpha + \lambda^2\gamma, \lambda\beta + \gamma, -\lambda\alpha - \gamma, \beta + \lambda^2\gamma) \tag{7}$$

whenever $\alpha^2 + \alpha\beta + \beta^2 = 3\lambda\gamma^2$ (Ramanujan's result was slightly pre-dated by a very similar general solution due to Schwering (see [3], p. 557).)

Despite these results, however, there is no known formula characterizing the **integral** solutions to equation (1). In this light, considering various families of solutions is of value. This paper categorizes and extends various families of solutions to equation (1). Many of the results may be found in Dickson [3] and Barbeau [1].

There are many other one-parameter families of solutions to equation (1) besides (2) and (3). Examples are:

$$\begin{aligned}
 (a, b, c, d) = & ((2x - 1)(2x^3 - 6x^2 - 1), (x + 1)(5x^3 - 9x^2 + 3x - 1), \\
 & 3x(x + 1)(x^2 - x + 1), 3x(2x - 1)(x^2 - x + 1));
 \end{aligned} \tag{8}$$

$$(a, b, c, d) = (x^3 + 1, 2x^3 - 1, x(x^3 - 2), x(x^3 + 1)); \tag{9}$$

$$(a, b, c, d) = (3x^2, 6x^2 \pm 3x + 1, 3x(3x^2 \pm 2x + 1), c + 1). \tag{10}$$

As before, one may let x be a rational number v/u and multiply through by an appropriate power of u to obtain a two-parameter family of **integral** solutions.

A strikingly dissimilar one-parameter family of solutions is due to Ramanujan. Letting

$$\begin{aligned}
 \frac{1 + 53x + 9x^2}{1 - 82x - 82x^2 + x^3} &= \sum_{n \geq 0} a_n x^n, \\
 \frac{2 - 26x - 12x^2}{1 - 82x - 82x^2 + x^3} &= \sum_{n \geq 0} b_n x^n, \\
 \frac{2 + 8x - 10x^2}{1 - 82x - 82x^2 + x^3} &= \sum_{n \geq 0} c_n x^n,
 \end{aligned}$$

yields

$$a_n^3 + b_n^3 = c_n^3 + (-1)^n.$$

This result produces "near misses" when considering Fermat's Last Theorem. Hirschhorn [4] has observed that Ramanujan's solutions are contained in

$$(a, b, c, d) = (u^2 + 7uv - 9v^2, -u^2 + 9uv + v^2, 2u^2 - 4uv + 12v^2, 2u^2 + 10v^2). \tag{11}$$

Some authors have given two-parameter families of solutions to equation (1) that could have been generated from a one-parameter solutions (as we have done earlier). Examples are:

$$(a, b, c, d) = (3u^2 + 5uv - 5v^2, 4u^2 - 4uv + 6v^2, 5u^2 - 5uv - 3v^2, 6u^2 - 4uv + 4v^2); \tag{12}$$

$$(a, b, c, d) = (3u^2 + 16uv - 7v^2, 6u^2 - 4uv + 14v^2, -3u^2 + 16uv + 7v^2, 6u^2 + 4uv + 14v^2). \quad (13)$$

Two-parameter solutions of (1) which do not arise from one-parameter solutions are not so plentiful. Ramanujan (see [1], pp. 35, 48) discovered

$$(a, b, c, d) = (u^7 - 3(v+1)u^4 + (3v^2 + 6v + 2)u, 2u^6 - 3(2v+1)u^3 + 3v^2 + 3v + 1, u^6 - 3v^2 - 3v - 1, u^7 - 3vu^4 + (3v^2 - 1)u). \quad (14)$$

In comparing the different families of solutions previously mentioned, one notices that the coefficients in the solution represented by (12) are the same as the values in equation (5). This generalizes to

Theorem 1: If

$$a^3 + b^3 + c^3 = d^3 \quad (15)$$

and

$$c(c^3 - a^2) = b(d^2 - b^2), \quad (16)$$

then

$$(ax^2 + cx - c)^3 + (bx^2 - bx + d)^3 + (cx^2 - cx - a)^3 = (dx^2 - bx + b)^3.$$

This theorem may be proved directly by expansion. It shows that a one-parameter family of solutions may sometimes be constructed from one solution. The next theorem shows exactly where Theorem 1 applies.

Theorem 2: The only solutions of equations (15)-(16) are:

(a) (trivial) solutions of the form $(a, b, c, d) = (a, b, -a, b)$;

(b) (scaled) solutions of the one-parameter system represented by (9), namely

$$(a, b, c, d) = (1 + u^3, u^4 - 2u, 2u^3 - 1, u^4 + u).$$

Proof: Substituting Euler's general solution (6) of (15) into (16) gives (after dividing by σ^3)

$$0 = 36\eta^2(\xi - \eta)(\xi^2 + 3\eta^2 - 1)(\xi^4 + 6\xi^2\eta^2 + \xi^2 + 9\eta^4 + 3\eta^2 + 1).$$

If $\eta = 0$ or $\xi^2 + 3\eta^2 - 1 = 0$, one falls into the first class of solutions. The only other possibility is if $\xi = \eta$, which yields

$$(a, b, c, d) = (8\eta^3 + 1, 16\eta^4 - 4\eta, 16\eta^3 - 1, 16\eta^4 + 2\eta).$$

Setting $u = 2\eta$ shows that this case falls into the second class of solutions. \square

Note that the second class of solutions is the same as (9). This solution is due to Vieta. Combining Theorems 1 and 2 generates a new two-parameter family of solutions to (1), namely

$$(a, b, c, d) = ((1 + u^3)x^2 + (2u^3 - 1)(x - 1), (u^4 - 2u)x(x - 1) + u^4 + u, (2u^3 - 1)x(x - 1) - u^3 - 1, (u^4 + u)x^2 - (u^4 - 2u)(x - 1)). \quad (17)$$

ACKNOWLEDGMENT

I would like to thank the anonymous referee for helpful comments which improved the presentation of this paper.

REFERENCES

1. E. J. Barbeau. *Power Play*. Washington, D.C.: MAA, 1997.
2. B. C. Berndt & S. Bhargava. "Ramanujan—for Lowbrows." *Amer. Math. Monthly* **100.7** (1993):644-656.
3. L. E. Dickson. *History of the Theory of Numbers*. Vol. 2. Washington, D. C.: Carnegie Institution of Washington, 1919-1923.
4. M. D. Hirschhorn. "An Amazing Identity of Ramanujan." *Math. Mag.* **68.3** (1995):199-201.
5. M. D. Hirschhorn. "A Proof in the Spirit of Zeilberger of An Amazing Identity of Ramanujan." *Math. Mag.* **69.4** (1996):267-69.
6. H. L. Keng. *Introduction to Number Theory*. New York: Springer, 1982.
7. Y. I. Perelman. *Algebra Can Be Fun*. Moscow: Mir Publishers, 1979.

AMS Classification Number: 11D25



NEW PROBLEM WEB SITE

Readers of *The Fibonacci Quarterly* will be pleased to know that many of its problems can now be searched electronically (at no charge) on the World Wide Web at

<http://problems.math.umr.edu>

Over 20,000 problems from 38 journals and 21 contests are referenced by the site, which was developed by Stanley Rabinowitz's MathPro Press. Ample hosting space for the site was generously provided by the Department of Mathematics and Statistics at the University of Missouri-Rolla, through Leon M. Hall, Chair.

Problem statements are included in most cases, along with proposers, solvers (whose solutions were published), and other relevant bibliographic information. Difficulty and subject matter vary widely; almost any mathematical topic can be found.

The site is being operated on a volunteer basis. Anyone who can donate journal issues or their time is encouraged to do so. For further information, write to:

Mr. Mark Bowron
Director of Operations, MathPro Press
P.O. Box 713
Westford, MA 01886 USA
bowron@my-deja.com