

# ODD FIBBINARY NUMBERS AND THE GOLDEN RATIO

LINUS LINDROOS, ANDREW SILLS, AND HUA WANG

ABSTRACT. The fibbinary numbers are positive integers whose binary representation contains no consecutive ones. We prove the following result: If the  $j$ th odd fibbinary is the  $n$ th odd fibbinary number, then  $j = \lfloor n\phi^2 \rfloor - 1$ .

## 1. BACKGROUND

The Fibonacci numbers  $\{F_n\}_{n \geq 0}$  are given by  $F_0 = 0$ ,  $F_1 = 1$ , and

$$F_n = F_{n-1} + F_{n-2}$$

for  $n \geq 2$ .

Recall the following theorem of Zeckendorf [1].

**Zeckendorf's Theorem.** *Every positive integer can be written uniquely as the sum of distinct, nonconsecutive Fibonacci numbers.*

The *Zeckendorf representation*  $z(n)$  of  $n \in \mathbb{N}$  is the unique  $k$ -tuple of decreasing nonconsecutive Fibonacci numbers whose sum is  $n$ . Note that although  $F_2 = F_1 = 1$ , we always associate 1 with  $F_2$  in the Zeckendorf representation.

For example,

$$z(4) = (3, 1) = (F_4, F_2),$$

$$z(5) = (5) = (F_5),$$

$$\text{and } z(100) = (89, 8, 3) = (F_{11}, F_6, F_4).$$

The sequence  $\{\text{fib}(n)\}_{n \geq 1}$  of *fibbinary numbers*<sup>1</sup> is given as follows: For  $n > 0$ , if  $z(n) = (F_{i_1}, F_{i_2}, \dots, F_{i_k})$  is the Zeckendorf representation of  $n$ , then

$$\text{fib}(n) := \sum_{j=1}^k 2^{i_j-2}.$$

For example,

$$\text{fib}(4) = \text{fib}(F_4 + F_2) = 2^{4-2} + 2^{2-2} = 101_2 = 5.$$

$$\text{fib}(5) = \text{fib}(F_5) = 2^{5-3} = 1000_2 = 8.$$

$$\text{fib}(100) = \text{fib}(F_{11} + F_6 + F_4) = 2^{11-2} + 2^{6-2} + 2^{4-2} = 1000010100_2 = 532,$$

where the 2 subscript indicates the usual binary (base two) representation.

---

The work of the third author was partially supported by a grant from the Simons Foundation (#245307).

<sup>1</sup>According to [3], the name “fibbinary” is due to Marc LeBrun: “. . . integers whose binary representation contains no consecutive ones and noticed that the number of such numbers with  $n$  bits was  $F_n$ .”—posting to `sci.math` by Bob Jenkins, July 17, 2002.

THE FIBONACCI QUARTERLY

2. STATEMENT OF MAIN RESULT RELATING ODD FIBBINARIES TO THE GOLDEN RATIO

It is easy to generate the odd fibbinary numbers from the binary representations and find the corresponding value of the original integer, e.g.,

- $1_2 = 2^0 = \text{fib}(F_2) = \text{fib}(1)$ ;
- $101_2 = 2^2 + 2^0 = \text{fib}(F_4 + F_2) = \text{fib}(4)$ ;
- $1001_2 = 2^3 + 2^0 = \text{fib}(F_5 + F_2) = \text{fib}(6)$ ;
- ...

Let  $\text{odfib}(n)$  denote the  $n$ th odd fibbinary number, i.e.,

$$\text{odfib}(1) = 1_2, \text{odfib}(2) = 101_2, \text{odfib}(3) = 1001_2, \dots$$

Let

$$Z(n) := \text{fib}^{-1}(\text{odfib}(n)),$$

so that

$$Z(1) = 1, Z(2) = 4, Z(3) = 6, Z(4) = 9, \dots$$

In other words, if the  $n$ th odd fibbinary number is the  $j$ th fibbinary number, then  $j = Z(n)$ .

This sequence

$$1, 4, 6, 9, 12, 14, 17, \dots$$

appears to be “A003622” in OEIS [2], defined as

$$\{\lfloor n\phi^2 \rfloor - 1\}_{n=1}^\infty = \{\lfloor n\phi \rfloor + n - 1\}_{n=1}^\infty \tag{1}$$

where

$$\phi = \frac{1 + \sqrt{5}}{2}$$

is the golden ratio, which satisfies  $\phi^2 = \phi + 1$ , and of course arises as

$$\lim_{n \rightarrow \infty} \frac{F_{n+1}}{F_n} = \phi.$$

$j$	$z(j)$	$\text{fib}(j)$	$\text{odfib}^{-1}(j)$
$1 = \lfloor \phi^2 \rfloor - 1 = Z(1)$	$(1) = (F_2)$	$1_2 = 1$	1
2	$(2) = (F_3)$	$10_2 = 2$	—
3	$(3) = (F_4)$	$100_2 = 4$	—
$4 = \lfloor 2\phi^2 \rfloor - 1 = Z(2)$	$(3, 1) = (F_4, F_2)$	$101_2 = 5$	2
5	$(5) = (F_5)$	$1000_2 = 8$	—
$6 = \lfloor 3\phi^2 \rfloor - 1 = Z(3)$	$(5, 1) = (F_5, F_2)$	$1001_2 = 9$	3
7	$(5, 2) = (F_5, F_3)$	$1010_2 = 10$	—
8	$(8) = (F_6)$	$10000_2 = 16$	—
$9 = \lfloor 4\phi^2 \rfloor - 1 = Z(4)$	$(8, 1) = (F_6, F_2)$	$10001_2 = 17$	4
10	$(8, 2) = (F_6, F_3)$	$10010_2 = 18$	—
11	$(8, 3) = (F_6, F_4)$	$10100_2 = 20$	—
$12 = \lfloor 5\phi^2 \rfloor - 1 = Z(5)$	$(8, 3, 1) = (F_6, F_4, F_2)$	$10101_2 = 21$	5

The correspondence displayed above is naturally conjectured to be true in general.

**Theorem 2.1.** *Let  $j$  be a positive integer such that the  $j$ th fibbinary number is odd. Suppose that this  $j$ th fibbinary number is the  $n$ th odd fibbinary number. Then*

$$j = Z(n) = \lfloor n\phi^2 \rfloor - 1 = \lfloor n\phi \rfloor + n - 1 \tag{2}$$

for any  $n \geq 1$ .

### 3. PROOF OF THEOREM 2.1

It is easy to check that (2) is true for small values of  $n$ . In the rest of this note we provide a general proof for  $n \geq 3$ .

Next, we record the following observation on  $\text{odfib}(n)$  as a lemma.

**Lemma 3.1.** *For any  $1 \leq k \leq F_{n-1}$  and  $n \geq 2$ , we have*

$$\text{odfib}(F_n + k) = 2^n + \text{odfib}(k).$$

*Proof.* The results follow since each digit in the binary representation of  $\text{odfib}(n)$  corresponds to a specific Fibonacci number.  $\square$

For example,

$$\text{odfib}(10) = \text{odfib}(8 + 2) = \text{odfib}(F_6 + 2) = 1000101_2 = 2^6 + 101_2 = 2^6 + \text{odfib}(2).$$

**Lemma 3.2.** *For any  $1 \leq k \leq F_{n-1}$  and  $n \geq 2$ , we have*

$$Z(F_n + k) = F_{n+2} + Z(k).$$

*Proof.* Since each digit in the binary representation of  $\text{odfib}(n)$  corresponds to a distinct Fibonacci number in the sum of  $Z(n)$  under the Zeckendorf representation, we can claim the same for  $Z(n)$ . That is,

$$\begin{aligned} Z(F_n + k) &= \text{fib}^{-1}(\text{odfib}(F_n + k)) \\ &= \text{fib}^{-1}(2^n + \text{odfib}(k)) \\ &= F_{n+2} + \text{fib}^{-1}(\text{odfib}(k)) \\ &= F_{n+2} + Z(k). \end{aligned}$$

$\square$

For example,

$$Z(10) = Z(F_6 + 2) = 25 = 21 + 4 = F_8 + Z(2).$$

From Lemma 3.2, we have

$$Z(F_n + k) - Z(F_n + k - 1) = Z(k) - Z(k - 1) \tag{3}$$

for  $2 \leq k \leq F_{n-1}$  and

$$Z(F_n + 1) = F_{n+2} + 1. \tag{4}$$

Now to show (2) for any  $n$ , we only need to show analogues of (3) and (4) for  $\lfloor n\phi^2 \rfloor - 1$ , i.e.,

$$\begin{aligned} \lfloor (F_n + k)\phi \rfloor - \lfloor (F_n + k - 1)\phi \rfloor &= \lfloor (F_n + k)\phi^2 \rfloor - \lfloor (F_n + k - 1)\phi^2 \rfloor \\ &= \lfloor k\phi^2 \rfloor - \lfloor (k - 1)\phi^2 \rfloor \\ &= \lfloor k\phi \rfloor - \lfloor (k - 1)\phi \rfloor \end{aligned} \tag{5}$$

for  $2 \leq k \leq F_{n-1}$ , and

$$\lfloor (F_n + 1)\phi \rfloor + F_n = \lfloor (F_n + 1)\phi^2 \rfloor - 1 = Z(F_n + 1). \tag{6}$$

**Remark.** Intuitively, this can be considered as using  $Z(F_n + 1)$  as the “stepping stone” to prove (2) for  $Z(F_n + k)$  for  $k = 2, 3, \dots, F_{n-1}$  using induction.

THE FIBONACCI QUARTERLY

In order to establish (5), it is essentially sufficient to show that  $\{F_n\phi\}$  is never large enough to affect the difference  $\lfloor k\phi \rfloor - \lfloor (k-1)\phi \rfloor$ , where  $\{x\} := x - \lfloor x \rfloor$ , the fractional part of the real number  $x$ .

We make use of the following fact, which is easily established by induction in  $n$ :

$$(-1)^n \tau^n = -F_n \tau + F_{n-1}, \tag{7}$$

where  $\tau = \frac{\sqrt{5}-1}{2} = \phi - 1$  satisfying  $\tau^2 = -\tau + 1$ . Consequently,

$$\{F_n \phi\} = \{F_n \tau\} = \{-(-\tau)^n\}$$

for any  $n$ .

Making use of the fact that  $\tau = \phi - 1$ , it suffices to show

$$\lfloor (F_n + k)\tau \rfloor - \lfloor (F_n + k - 1)\tau \rfloor = \lfloor k\tau \rfloor - \lfloor (k-1)\tau \rfloor. \tag{8}$$

To show (8), simply consider

$$\{k\tau\} \pm \{F_n\tau\}$$

for any  $1 \leq k \leq F_{n-1}$ . We will show that this value never reaches 1 or goes below zero and hence,  $\{F_n\tau\}$  will not affect  $\lfloor k\tau \rfloor - \lfloor (k-1)\tau \rfloor$ .

(i) To show  $\{k\tau\} + \{F_n\tau\} < 1$ , consider the Zeckendorf representation of  $k$  as the sum of nonconsecutive Fibonacci numbers.

If  $k < F_{n-1}$ , then

$$k = F_{a_1} + F_{a_2} + \dots + F_{a_s}$$

where

$$1 \leq a_1 \leq a_2 - 2 \leq a_2 \leq a_3 - 2 \leq \dots \leq a_s \leq n - 2.$$

Then

$$\begin{aligned} \{k\tau\} + \{F_n\tau\} &= \{(F_{a_1} + \dots + F_{a_s})\tau\} + \{F_n\tau\} \\ &\leq \tau^{a_1} + \tau^{a_2} + \dots + \tau^{a_s} + \tau^n \\ &< \tau + \tau^3 + \tau^5 + \dots \\ &= \frac{\tau}{1 - \tau^2} \\ &= 1. \end{aligned}$$

If  $k = F_{n-1}$ , we have

$$\{k\tau\} + \{F_n\tau\} = \tau^{n-1} + \tau^n \leq \tau^2 + \tau^3 < 1$$

for any  $n \geq 3$ .

(ii) To show  $\{k\tau\} - \{F_n\tau\} > 0$ , simply note that

$$\begin{aligned} \{k\tau\} - \{F_n\tau\} &= \{(F_{a_1} + \dots + F_{a_s})\tau\} - \{F_n\tau\} \\ &\geq \tau^{a_1} - \tau^{a_2} - \dots - \tau^{a_s} - \tau^n \\ &> \tau^{a_1} - \tau^{a_1+2} - \tau^{a_1+4} - \dots \\ &= \tau^{a_1} \left( 1 - \frac{\tau^2}{1 - \tau^2} \right) \\ &> 0 \end{aligned}$$

if  $k < F_{n-1}$  and

$$\{k\tau\} - \{F_n\tau\} = \tau^{n-1} - \tau^n > 0$$

if  $k = F_{n-1}$ .

## ODD FIBBINARY NUMBERS AND THE GOLDEN RATIO

Cases (i) and (ii) imply that

$$\lfloor F_n \tau + k\tau \rfloor = \lfloor F_n \tau \rfloor + \lfloor k\tau \rfloor.$$

Thus (8) and (5) are proved.

By (4), (6) is equivalent to

$$\lfloor (F_n + 1)\phi \rfloor = F_{n+1} + 1. \tag{9}$$

Using (7) it follows that

$$\begin{aligned} \lfloor (F_n + 1)\phi \rfloor &= F_n + 1 + \lfloor (F_n + 1)\tau \rfloor \\ &= F_n + 1 + \lfloor F_{n-1} - (-\tau)^n + \tau \rfloor \\ &= F_n + 1 + F_{n-1} + \lfloor \tau - (-\tau)^n \rfloor \\ &= F_{n+1} + 1 \end{aligned}$$

for  $n \geq 3$ . Therefore, (9) is true and hence, (6) is proved. □

### ACKNOWLEDGEMENT

We thank the anonymous referee for carefully reading the manuscript and providing helpful suggestions.

### REFERENCES

- [1] E. Zeckendorf, *Représentation des nombres naturels par une somme des nombres de Fibonacci ou de nombres de Lucas*, Soc. Roy. Sci. Liège, **41** (1972), 179–182.
- [2] OEIS Foundation Inc. (2011), The On-Line Encyclopedia of Integer Sequences, <http://oeis.org/A003622>.
- [3] OEIS Foundation Inc. (2011), The On-Line Encyclopedia of Integer Sequences, <http://oeis.org/A003714>.

MSC2010: 11B39

DEPARTMENT OF MATHEMATICAL SCIENCES, GEORGIA SOUTHERN UNIVERSITY, STATESBORO, GA 30460  
*E-mail address:* [1101383@georgiasouthern.edu](mailto:1101383@georgiasouthern.edu)

DEPARTMENT OF MATHEMATICAL SCIENCES, GEORGIA SOUTHERN UNIVERSITY, STATESBORO, GA 30460  
*E-mail address:* [asills@georgiasouthern.edu](mailto:asills@georgiasouthern.edu)

DEPARTMENT OF MATHEMATICAL SCIENCES, GEORGIA SOUTHERN UNIVERSITY, STATESBORO, GA 30460  
*E-mail address:* [hwang@georgiasouthern.edu](mailto:hwang@georgiasouthern.edu)