

Equation (4) is of the same form as (1)' with k replaced by $k + 1$. Hence, etc.

Similarly other results can be proved for all positive integral values of n .

REFERENCES

1. A. F. Horadam, "Basic Properties of a Certain Generalized Sequence of Numbers," The Fibonacci Quarterly, Vol. 3, 1965, pp. 161-176.
2. A. F. Horadam, "Complex Fibonacci Numbers and Fibonacci Quaternions," Amer. Math. Monthly, Vol. 70, 1963, pp. 289-291.
3. K. Subba Rao, "Some Properties of Fibonacci Numbers," Amer. Math. Monthly, Vol. 60 (1953), pp. 680-684.

[Continued from p. 91.]

3. Tobias Dantzig, Bequest of the Greeks, New York, Charles Scribner's Sons, 1955, pp. 61-62.
4. Marvin H. Holt, "Mystery Puzzler and Phi," The Fibonacci Quarterly, Vol. 3, No. 2, April 1965, pp. 135-138.
5. H. E. Huntley, "The Golden Cuboid," The Fibonacci Quarterly, Vol. 2, No. 3, October 1964, p. 184⁺.
6. Joseph A. Raab, "A Generalization of the Connection Between the Fibonacci Sequence and Pascal's Triangle," The Fibonacci Quarterly, Vol. 1, No. 3, October 1963, pp. 21-31.
