Theorem 2 is a dual to Theorem 1 and corresponds to the dual of the Zeckendorf theorem for Fibonacci numbers  $\lceil 4 \rceil$ .

## REFERENCES

- 1. J. L. Brown, Jr., "Note on Complete Sequences of Integers," American Mathematical Monthly, Vol. 68, No. 6, June-July, 1961, pp. 557-560.
- 2. C. G. Lekkerkerker, "Voorstelling van natuurlijke getallen door een som van getallen van Fibonacci," Simon Stevin, Vol. 29, 1951-52, pp. 190-195.
- 3. J. L. Brown, Jr., "Zeckendorf's Theorem and Some Applications," The Fibonacci Quarterly, Vol. 2, No. 3, October, 1964, pp. 163-168.
- 4. J. L. Brown, Jr., "A New Characterization of Fibonacci Numbers," The Fibonacci Quarterly, Vol. 3, No. 1, February 1965, pp. 1-8.

## ASSOCIATION MEETING

The Fibonacci Association held its Fall Meeting on October 18th at San Jose State College. Following was the Program:

## MORNING SESSION

9:30 a.m.	SOCIAL GATHERING
10:00 — 10:45	TEST FOR THE PRIMALITY OF MERSENNE NUMBERS Douglas Lind, Stanford University
10:45 — 11:30	WEB SEQUENCES George Ledin, Jr., University of San Francisco
11:30 — 12 Noon	OPPORTUNITY FOR GENERAL DISCUSSION

## AFTERNOON SESSION

1:15 — 2:00	FIBONACCI AND RELATED SERIES IN COMBINATORICS Prof. D. H. Lehmer, University of Calif., Berkeley
2:00 — 2:45	MARKOV-FIBONACCI RELATIONS Prof. Gene Gale, San Jose State College
2:45 — 3:30	IT'S GENERALIZED! WHAT'S NEXT? Prof. V. C. Harris, San Diego State College

\* \* \* \* \*