



Fibonacci-18

Conference Program

The Eighteenth International Conference on
Fibonacci Numbers and Their Applications

Halifax, Nova Scotia, Canada

July 1 – 8, 2018



DALHOUSIE 1818
UNIVERSITY 2018

Contents

Welcome	5
Conference Schedule	7
Social Activities	16
Abstracts	17
The Paul Bruckman Prizes	35
The Fibonacci Portrait	36
Food	37
Books	39
Maps	43

For **wireless internet** access on campus, either

- connect to **eduroam** and use your eduroam credentials;
- or ask Karl Dilcher for a temporary login and connect to **Dal-WPA2**.

**Dalhousie University is located in Mi'kma'ki, the
ancestral and unceded territory of the Mi'kmaq.**

Welcome

Welcome to Dalhousie University, to Halifax, and for many of you, to Canada! This conference begins as we celebrate Canada Day (July 1st), with Monday, July 2nd a national holiday this year. Canada celebrates the 151st anniversary of Confederation, an act that took place in Charlottetown in the neighbouring small province of Prince Edward Island.

We are now also exactly half-way through Dalhousie University's bi-centennial year, and this Fibonacci Conference is an official part of the scientific events that mark this significant anniversary. It is perhaps appropriate that the 18th International Conference on Fibonacci Numbers and Their Applications takes place in '18, at a university that was founded in 1818. Needless to say, 18 is not a Fibonacci number, but it is a Lucas number, and thus connected with the second-most important name attached to this conference.

Some words of thanks are now in order. We received financial support for this conference from AARMS (the Atlantic Association for Research in the Mathematical Sciences) and from the Faculty of Science of Dalhousie University. I would also like to acknowledge the support of the Department of Mathematics and Statistics and its office staff, and finally I'd like to thank Asmita Sodhi, Mason Maxwell, and especially Keith Johnson for their help and support.

I thank you all for coming, and I wish everybody a very pleasant stay in Halifax, and an interesting and successful conference.

Karl Dilcher.



Conference Schedule

All talks take place in the “Scotiabank Auditorium” of the Marion McCain Arts & Social Sciences Building.

Monday, July 2

Morning Session

8:15	Registration opens
8:45	Opening Remarks
9:00–9:25	Christian Ballot: <i>Variations on Catalan Lucasnomials</i>
9:30–9:55	Heiko Harborth: <i>A Conjecture for Pascal’s Triangle</i>
10:00–10:30	— Coffee Break —
10:30–10:55	William Webb: <i>What Makes A “Nice” Identity?</i>
11:00–11:25	Arthur T. Benjamin: <i>Some Bingo Paradoxes</i>
11:30–11:55	Steven Miller: <i>From Monovariants to Zeckendorf Decompositions and Games</i>
12:00–2:00	— Lunch Break —
Note:	For members of the Fibonacci Association Board:
12:00–2:00	Board Meeting & Luncheon Dalhousie University Club

Monday, July 2

Afternoon Session

2:00–2:25 Sadjia Abbad:

Companion Sequences Associated to the r -Fibonacci Sequence

2:30–2:55 Paul Young:

The Power of 2 Dividing a Generalized Fibonacci Number

3:00–3:30 — Coffee Break —

3:30–3:55 Antara Mukherjee:

The Geometric Interpretation of Some Fibonacci Identities in the Hosoya Triangle

4:00–4:25 J. C. Saunders:

On (a, b) Pairs in Random Fibonacci Sequences

4:30–4:55 Marc Chamberland:

Arctan Formulas and π

Tuesday, July 3

Morning Session

9:00–9:25	Orli Herscovici: <i>New Degenerated Bernoulli and Euler Polynomials Arising from Non-Classical Umbral Calculus</i>
9:30–9:55	Lin Jiu: <i>Bessel Random Walks and Identities for Higher-Order Bernoulli and Euler Polynomials</i>
10:00–10:10	— Group Photo — Location to be announced —
10:10–10:30	— Coffee Break —
10:30–10:55	Sam Northshield: <i>Re³ counting the Rationals</i>
11:00–11:25	Larry Ericksen: <i>Properties of Polynomials that Encode Representations</i>
11:30–11:55	Paul K. Stockmeyer: <i>Discovering Fibonacci Numbers, Fibonacci Words, and a Fibonacci Fractal in the Tower of Hanoi</i>
12:00–1:30	— Lunch Break —

Tuesday, July 3

Afternoon Session

1:30–1:55 Susanna Spektor:

On a ψ_1 -Norm Estimate of Sums of Dependent Random Variables Using Simple Random Walks on Graphs

2:00–2:25 Meliza Contreras González:

Counting Independent Sets on Bipolygonal Graphs

2:30–2:55 Thotsaporn ‘Aek’ Thanatipanonda:

Statistics of Domino Tilings on a Rectangular Board

3:00–3:30 — Coffee Break —

3:30–3:55 Elif Tan:

A Note on Conditional Divisibility Sequences

4:00–4:25 Tanay Wakhare:

Structural Identities for Multiple Zeta Values

4:30–4:55 Paul Young:

Global Series for Zeta Functions

Wednesday, July 4

Morning Session

9:00–9:55	The Édouard Lucas Memorial Lecture Hugh C. Williams: <i>Mersenne, Fibonacci and Lucas: The Mersenne Prime Story and Beyond</i>
10:00–10:30	— Coffee Break —
10:30–10:55	Burghard Herrmann: <i>How Integer Sequences Find Their Way Into Areas Outside “Pure Mathematics”</i>
11:00–11:25	Dale Gerdemann: <i>Images From Zeckendorf and Other Numerical Representations</i>
11:30–11:55	Bruce Boman: <i>Geometric Branching Patterns Based on the p-Fibonacci Numbers: Self-Similarity Across Different Degrees of Branching and Multiple Dimensions</i>
12:00–1:30	— Lunch Break —
1:30	Afternoon Excursion

Thursday, July 5

Morning Session

9:00–9:25	Peter Anderson: <i>More Remarkable Continued Functions</i>
9:30–9:55	Bir Kafle: <i>Pell Numbers of the Form $2^a + 3^b + 5^c$</i>
10:00–10:30	— Coffee Break —
10:30–10:55	Karyn McLellan: <i>A Problem on Generating Sets Containing Fibonacci Numbers</i>
11:00–11:25	Curtis Cooper: <i>Some Generalized High Order Fibonacci Identities</i>
11:30–11:55	Scott Cameron: <i>A Linear Algebra Problem Related to Legendre Polynomials</i>
12:00–1:30	— Lunch Break —

Thursday, July 5

Afternoon Session

1:30–1:55 Steven Edwards:

Generalizations of Delannoy Numbers

2:00–2:25 Barry Balof:

Selfish Sets, Posets, Tilings and Bijections

2:30–2:55 Michael Allen:

*A New Combinatorial Interpretation of the
Fibonacci Numbers Squared*

3:00–3:30 — Coffee Break —

3:30–3:55 Russell Jay Hendel:

*Proof and Formulation of a Tagiuri-Generating-Method
Conjecture*

4:00–4:25 Bob Bastasz:

Digital Loop Systems

Friday, July 6

Morning Session

9:00–9:25	Clark Kimberling: <i>Linear Complementary Equations and Systems</i>
9:30–9:55	Abdullah Al-Shaghay: <i>Irreducibility and Roots of a Class of Polynomials</i>
10:00–10:30	— Coffee Break —
10:30–10:55	William Webb: <i>Proving Identities In Arbitrary Fields</i>
11:00–11:25	Osman Yürekli: <i>A Pascal-like Triangle From a Special Function</i>
11:30–11:55	Burghard Herrmann: <i>Visibility in a Pure Model of Golden Spiral Phyllotaxis</i>
12:00–1:30	— Lunch Break —

Friday, July 6

Afternoon Session

1:30–1:55	Prapanpong Pongsriam: <i>Fibonacci and Lucas Numbers Which Have Exactly Three Prime Factors and Some Unique Properties of F_{18} and L_{18}</i>
2:00–2:25	Christophe Vignat: <i>Finite Generating Functions for the Sum-of-Digits Sequence</i>
2:30–2:55	Kouichi Nakagawa: <i>Exact Periodicity of Generalized Fibonacci and Tribonacci Sequences</i>
3:00–3:30	— Coffee Break —
3:30–4:30	Clark Kimberling (Coordinator): <i>Problem Session</i>
4:30	Closing Remarks

Social Activities

Sunday, July 1:

4:00–9:00 pm: Registration desk is open – conference material can be picked up.

7:00–9:00 pm: Welcoming Reception.

Both in the lobby of the Marion McCain Arts & Social Sciences Building.

Tuesday, July 3:

10:00 am: Group photo; location to be announced.

Wednesday, July 4:

1:30 pm: Beginning of afternoon excursion to Peggy's Cove and Lunenburg.

Bus pick-up on Campus; exact location to be announced.

Thursday, July 5:

Conference Dinner in Hubbards (around 45 Minutes from Halifax).

Locations and time for bus pick-up to be announced.

Saturday, July 7:

Optional full-day trip to Cape Split with hike to the tip of the peninsula there. The hike is relatively easy, and is 1 1/2 hours each way from the parking lot. The trip to Cape Split also offers some scenic views along the way. Transportation will be with private vehicles or a rented van, depending on the number of participants. There will be a sign-up sheet during the conference.

Abstracts

The Édouard Lucas Memorial Lecture

Hugh C. Williams, University of Calgary, Calgary, AB, Canada

Mersenne, Fibonacci and Lucas:

The Mersenne Prime Story and Beyond

On Dec. 26 of last year, it was announced that the 50th known Mersenne prime had been identified. This is an enormous number of 23,249,425 decimal digits and is currently the largest known prime number. In spite of the size of this number we are able to prove it prime by a simple algorithm that was discovered in 1876 by Édouard Lucas. Lucas discovered this procedure as a result of his examination of the properties of Fibonacci numbers.

In this talk I will briefly discuss the development of the concept of a Mersenne prime and then describe Lucas' ideas concerning how the primality of such numbers can be established. I will also detail some aspects of Lucas' career and conclude with a discussion of his unsuccessful search for a generalization of his technique.

Abstracts – Contributed Talks

Abstracts are listed in alphabetical order by speaker.

Sadjia Abbad, Saad Dahlab University, Blida, Algeria

Companion Sequences Associated to the r -Fibonacci Sequence

In this talk, we define the r -Lucas sequences of type s . These sequences constitute a family of companion sequences of the generalized r -Fibonacci sequences. We establish the corresponding Binet formula and evaluate generating functions. Therefore we extend the definition of $V_n^{(r,s)}$ to negative n . Also, we exhibit some convolution relations which generalize some known identities such as Cassinis.

(Joint work with Hacène Belbachir.)

Michael A. Allen, Mahidol University, Bangkok, Thailand

A New Combinatorial Interpretation of the Fibonacci Numbers Squared

We consider the tiling of an n -board (a $1 \times n$ array of square cells of unit width) with half-squares ($\frac{1}{2} \times 1$ tiles) and $(\frac{1}{2}, \frac{1}{2})$ -fence tiles. A $(\frac{1}{2}, \frac{1}{2})$ -fence tile is composed of two half-squares separated by a gap of width $\frac{1}{2}$. We show that the number of ways to tile an n -board using these types of tiles equals F_{n+1}^2 where F_n is the n th Fibonacci number. We use these tilings to devise combinatorial proofs of identities relating the Fibonacci numbers squared to one another. Some of these identities appear to be new.

Abdullah Al-Shaghay, Dalhousie University, Halifax, NS, Canada

Irreducibility and Roots of a Class of Polynomials

In 2012 Harrington studied the factorization of trinomials of the form $x^n + cx^{n-1} + d \in \mathbb{Z}[x]$. As an application of these results on trinomials, he proves factorization properties of polynomials of the form $x^n + c(x^{n-1} + \dots + x + 1) \in \mathbb{Z}[x]$. In this presentation, results regarding the factorization and roots of polynomials of the form $x^n + c(x^{n-a-1} + \dots + x + 1) \in \mathbb{Z}[x]$ are introduced. Analogously to Harrington, quadrinomials of the form $x^{n+1} \pm x^n \pm cx^{n-a} \pm c$ associated to our polynomials are considered.

Peter G. Anderson, RIT, Rochester, NY, USA

More Remarkable Continued Functions

Some sequences of linear functions of the form $T_n = \frac{x+b_n}{c_n}$ which satisfies a Fibonacci-like composition rule, $T_{n+1} = T_n \circ T_{n-1}$ has a sequence of fixed points $f_n = b_n/(c_n - 1)$ involving a remarkable continued fraction $[a_0, a_1, \dots]$ (either finite for any T_n or infinite for $\lim_{n \rightarrow \infty} T_n$) satisfying a Fibonacci-like multiplication rule, $a_{m+1} = a_m a_{m-1}$, for $m \geq 1$.

For every pair of positive integers, a_0, a_1 , there is a pair of functions, T_0, T_1 , giving the sequence T_n , as above, with fixed points involving the continued fraction as described above.

Christian Ballot, University of Caen, Caen, France

Variations on Catalan Lucasnomials

If $U = (U_n)_{n \geq 0}$ is a sequence of nonzero integers, then one may consider the generalized binomial coefficients, $\binom{m}{n}_U$, with respect to U . They are defined for $m \geq n \geq 0$ as follows

$$\binom{m}{n}_U = \frac{U_m U_{m-1} \dots U_{m-n+1}}{U_n U_{n-1} \dots U_1},$$

if $m \geq n \geq 1$, and as 1, if $n = 0$.

We will solely concentrate on the case when $U = U(P, Q)$ is a non-degenerate fundamental Lucas sequence, i.e., a second-order linear recurrent sequence with $U_0 = 0$, $U_1 = 1$ and $U_n \neq 0$ for $n \geq 2$ which satisfies

$$U_{n+2} = PU_{n+1} - QU_n, \text{ for all } n \geq 0,$$

where P and Q are nonzero integers. Those generalized binomial coefficients are referred to as Lucasnomials. Thus, the sequence $I = (I_n)_{n \geq 0}$ of natural numbers is the particular fundamental Lucas sequence $U(2, 1)$ which yields the ordinary binomial coefficients.

We know $n + 1$ divides $\binom{2n}{n}$ for all $n \geq 1$. If k is an integer not 1, then there are infinitely many $n \geq 1$ for which $n + k$ does not divide $\binom{2n}{n}$. As it happens this Catalan phenomenon remains true, or nearly so, for Lucasnomials. That is, for $\gcd(P, Q) = 1$,

$$\frac{1}{U_{n+k}} \binom{2n}{n}_U$$

is an integer for all $n \geq 1$ iff $k = 1$, or $k = 2$ and $U = U(1, 2)$. We will review further extensions of these results, in particular with respect to Lucasnomial Fuss-Catalan numbers and go in detail over theorems surrounding this phenomenon. Open problems will be outlined.

Barry Balof, Whitman College, Walla Walla, WA, USA
Selfish Sets, Posets, Tilings and Bijections

A subset of the integers $\{1, 2, \dots, n\}$ is *selfish* if it contains its own cardinality as an element. Those sets for which the minimal element is the cardinality (referred to by Grimaldi as *extraordinary* sets) are enumerated by the Fibonacci Numbers. In a 2013 paper, Grimaldi and Rickert introduced a partial order on these extraordinary sets. In this talk, we will establish natural bijections between the subsets and domino-square tilings to give a new interpretation to some combinatorial identities.

Bob Bastasz, Missoula, MT, USA
Digital Loop Systems

A digital loop system $S[m, l]$ is a set of periodic sequences based on a l -order linear recurrence in a finite field \mathbb{F}_m . Each sequence, called a loop, is expressed as a Lyndon word consisting of the digits in its least period and can be uniquely specified by a minimal element, which is a l -tuple pre-necklace. The periods of all distinct loops in a system sum to m^l . For example, the Fibonacci sequence (mod 10), with a period of 60, is one of six digital loops contained the system $S[10, 2]$, whose periods sum to 10^2 .

A basic property of a digital loop system is the number of distinct loop periods, c . Of particular interest are systems in which m is a prime and c is two or three. If $S[m, l]$ has $c=2$, it is proposed that $S[m^i, l]$ has $c = i + 1$ where i is a positive integer. Cases where the same period can be found for loops in $S[m, l]$ and $S[m^2, l]$ will be discussed.

Arthur T. Benjamin, Harvey Mudd College, Claremont, CA, USA
Some Bingo Paradoxes

In the game of Bingo, when many cards are in play, it is much more probable that the winning card is horizontal than vertical. We will explore this and other paradoxes. Fibonacci numbers and q -binomial coefficients make a brief appearance.

Bruce M. Boman, University of Delaware, Newark, DE, USA
Geometric Branching Patterns Based on the p -Fibonacci Numbers: Self-similarity Across Different Degrees of Branching and Multiple Dimensions

Branching patterns occur throughout nature and are often described by the Fibonacci numbers. While the regularity of these branching

patterns in biology can be described by the Fibonacci numbers, the branches (leaves, petals, offshoots, limbs, etc) are often variegated (size, color, shape, etc). To begin to understand how these patterns arise, we considered different branching patterns based on the p -Fibonacci numbers. In our model, different branch patterns were created based on a specific number of decreasing-sized branches that arise from a main branch (termed the degree of branching). It was assumed that the ratio between the sizes of pairs of consecutive branches (ordered by size) equals the ratio of the largest branch size to the sum of the largest and smallest branch sizes. Generation of these branching structures illustrates that pattern self-similarities occur across different degrees of branching and multiple dimensions. Conclusion: studying geometric branching patterns based on the p -Fibonacci numbers begins to show how the regularity in branching patterns might occur in biology.

(Joint work with Gilberto Schleiniger).

Scott Cameron, Dalhousie University, Halifax, NS, Canada

A Linear Algebra Problem Related to Legendre Polynomials

I introduce a problem which piqued my interest, namely a question asked in the context of simple linear algebra, and then generalize this problem to investigate further properties. This leads to a study of families of polynomial coefficients for kernels of shifted Legendre polynomials, and the properties which they have. It turns out that there is a general formula for the generating function of each of these families.

Marc Chamberland, Grinnell College, Grinnell, IA, USA

Arctan Formulas and π

There are many interesting formulas connected to the arctan function. For centuries, Machin-like formulas, such as

$$\frac{\pi}{4} = 4 \arctan \left(\frac{1}{5} \right) - 4 \arctan \left(\frac{1}{239} \right)$$

were the main technique used for calculating Pi. Starting with a geometric motivation, we build several new arctan formulas, for example,

$$\begin{aligned} \pi = & \arctan \left(a \sqrt{\frac{a+b+c}{abc}} \right) + \arctan \left(b \sqrt{\frac{a+b+c}{abc}} \right) \\ & + \arctan \left(c \sqrt{\frac{a+b+c}{abc}} \right) \end{aligned}$$

when $a, b, c > 0$. (Joint work with Eugene Herman.)

Meliza Contreras González, Universidad Autónoma de Puebla,
Puebla, Mexico

Counting Independent Sets on Bipolygonal Graphs

We consider the sequence $\beta_{s,k} = F_s \cdot F_{k-s}$ for $k > 0, 1 \leq s \leq k - 1$, introduced in [1], that is formed by the product of two Fibonacci numbers with complementary indexes. The values of this sequence allow us to compute the number of independent sets on bipolygonal graphs, which are graphs formed by two polygons C_i and C_j joined by an edge $e = \{x, y\}$, with $x \in V(C_i)$ and $y \in V(C_j)$. We denote this class of graphs as $H_{i,j}$. In particular, when the polygons C_i and C_j are hexagons, then $H_{i,j}$ is the primitive graph used to form chains of polyphenylene compounds [2].

We apply the edge division rule to decompose $H_{i,j}$ and to use the values in the sequence $\beta_{s,k}$ for computing the number of independent sets of $H_{i,j}$, denoted as $i(H_{i,j})$. In fact, $i(H_{i,j}) = F_{i+1} \cdot F_{j+1} + F_{i+1} \cdot F_{j-1} + F_{i-1} \cdot F_{j+1}$. Fixing $k \geq 6$, and $k = i + j$, we consider the different subgraphs formed by the variations: $3 \leq i, j \leq (k - 3)$. We analyze all possible size combinations for C_i and C_j , fixing $i + j$ as a constant k .

In addition, the way to compute $H_{i,j}$ allow us to determine extremal topologies for $i(H_{i,j})$. The extremal values are identified when the greatest variation (entropy) between the sizes of the polygons C_i and C_j is achieved. The minimum value corresponds to $|C_j| - |C_i| = 6$, and the maximum value is given when $|C_j| - |C_i| = 4$.

(Joint work with Guillermo De Ita Luna and Pedro Bello López.)

References

- [1] G. De Ita, J. R. Marcial, J. A. Hernández, R. M. Valdovino, Extending Extremal Polygonal Arrays for the Merrifield-Simmons Index, *Lecture Notes in Computer Science*, **10267** (2017), 22–31.
 [2] Došlić T., Litz M. S., Matchings and Independent Sets in Polyphenylene Chains, *MATCH Commun. Math. Comput. Chem.*, **67** (2012), 313–330.
-

Curtis Cooper, Univ. of Central Missouri, Warrensburg, MO, USA
Some Generalized High Order Fibonacci Identities

The Gelin-Cesáro identity states that for integers $n \geq 2$,

$$F_{n-2}F_{n-1}F_{n+1}F_{n+2} - F_n^4 = -1,$$

where F_n denotes the n th Fibonacci number. Horadam generalized the Fibonacci sequence by defining the sequence W_n where $W_0 = a$, $W_1 = b$, and $W_n = pW_{n-1} - qW_{n-2}$ for $n \geq 2$ and a , b , p and q are integers

and $q \neq 0$. Using this sequence, Melham and Shannon generalized the Gelin-Cesáro identity by proving that for integers $n \geq 2$,

$$W_{n-2}W_{n-1}W_{n+1}W_{n+2} - W_n^4 = eq^{n-2}(p^2 + q)W_n^2 + e^2q^{2n-3}p^2,$$

where $e = pab - qa^2 - b^2$. We will discover and prove some similar generalized high order Fibonacci identities.

Steven Edwards, Kennesaw State University, Marietta, GA, USA
Generalizations of Delannoy Numbers

The Delannoy numbers $D(m, n)$ count the number of lattice paths from $(0, 0)$ to (m, n) , where the allowable steps are up, right, and diagonal. The Delannoy numbers satisfy the recursion $D(m, n) = D(m, n - 1) + D(m - 1, n) + D(m - 1, n - 1)$. By restricting the number of diagonal steps allowed, we construct collections of generalized Delannoy numbers. The generalized Delannoy numbers satisfy the same recursion as the Delannoy numbers. There are many relations relating these numbers, and these relations, in turn, produce binomial identities which generalize known identities. Some of these identities provide insights into intrinsic properties of Pascal's triangle.

(Joint work with William Griffiths.)

Larry Ericksen, Millville, NJ, USA
Properties of Polynomials that Encode Representations

We present hyperbinary properties of two-variable Stern polynomials, with extension to hyper b-ary representations of integers. Continued fractions are also constructed from polynomial analogues of Lucas sequences.

(Joint work with Karl Dilcher.)

Dale Gerdemann
Images from Zeckendorf and Other Numerical Representations

Images, many of them fractal, can be generated from generalizations of numerical representations, primarily Golden Ratio Base, Zeckendorf and a variant of Zeckendorf which uses negatively indexed Fibonacci numbers (due to Martin Bunder). Bit sequences can be extracted from these representations that can be used to guide a walk in the plane, using color coding to represent the number of times each lattice point is encountered. Several algorithms can be used to convert the binary bit-pattern sequence into directional information, but all rely on the use of a division test using a prespecified divisor. The test may be on a count

of the total number of steps already taken in the walk or it may be on a more restricted count of just the steps corresponding to a 1-bit in the bit sequence. The idea is extended to Lucas numbers of the first kind: $U_0 = 0$, $U_1 = 1$, $U_n = sU_{n-1} + tU_{n-2}$. A variety of examples for various s, t and divisor can be seen at <https://bit.ly/2KcKSL7>. Fractal images exhibit symmetry at specific points in their construction. Based on experimentation, these points may correspond to Lucas numbers of the first or second kind. Possibly path lengths required to construct an image of some sort could be used to provide a counting interpretation of other combinatorial integer sequences. To test this idea, images are constructed for OEIS A181926 (diagonal sums of Fibonomial triangle).

Heiko Harborth, TU Braunschweig, Braunschweig, Germany
A Conjecture for Pascal's Triangle

For a prime number p consider Pascal's triangle reduced modulo p . Let $a_i(n)$ count the number of residues i in row n . If the linear combinations

$$c_0 a_0(n) + c_1 a_1(n) + \dots + c_{p-1} a_{p-1}(n) = 0$$

are fulfilled then it is conjectured by H.-D. Gronau and M. Krueppel that $c_i = 0$ for $0 \leq i \leq p - 1$. Partial results are presented.

Russell Jay Hendel, Towson University, Towson, MD, USA
Proof and Formulation of a Tagiuri-Generating-Method Conjecture

The Tagiuri Generating Method (TGM) generates one-parameter, infinite, families, of Fibonacci identities. To describe, $I(q)$, the q -th member of a family of identities, we need functions $s_p(q)$, $s_n(q)$, $s(q)$, and $m(q)$. Let $P = \prod_{j=1}^{m(q)} F_{n+a_j}$ with the a_j , $1 \leq j \leq m(q)$, parameters. TGM *starts* with an identity of the form $(s_p(q) - s_n(q))P = s_p(q)P - s_n(q)P$. TGM then requires replacement of one product $F_{n+a_k}F_{n+a_l}$ in each of the $s(q) = s_p(q) + s_n(q)$ summands on the right-hand side of the *start* identity with the corresponding right-hand side of the basic Tagiuri identity $F_{n+a_k}F_{n+a_l} = F_n F_{n+a_k+a_l} + (-1)^n F_{a_k} F_{a_l}$. Since the start and Tagiuri identities are true, $I(q)$ is also true. In practice, we map $\{a_j, 1 \leq j \leq m(q)\}$ to a subset of the integers symmetric around 0 (and excluding 0 if $m(q)$ is even). Particular examples of TGM families have been explored in FQ articles and conferences (Fibonacci (CAEN), West Coast Number Theory, MASON II). Main results are typically expressed as statements about the $I(q)$ -*index histograms*, $H_q = \{(x, c_q(x)) : x \in \mathbb{Z}\}$, where for each integer x , $c_q(x)$

counts the number of occurrences of F_{n+x} in $I(q)$. In this talk, we prove, *Under mild restrictions, $\#\{c_q(x) : (x, c_q(x)) \in H_q, x, q \in \mathbb{Z}, q \geq 1\} \leq c$, that is, for each $q \geq 1$, the number (cardinality, $\#$) of distinct index-counts in $I(q)$ is bounded above by a very small computable constant, c , independent of q .* The theorem is proven by presenting a single proof unifying all previous examples. The presentation closes by reviewing the history of Fibonacci-Lucas identities and showing a very recent trend to studying families of identities instead of individual identities or proof methods.

Burghard Herrmann, Köln, Germany

How Integer Sequences Find Their Way Into Areas Outside “Pure Mathematics”

Integers are considered on the surface of a cylinder as in the model of the pineapple in Coxeter [Introduction to Geometry, Chapter 11.5 Phyllotaxis (literally “leaf arrangement”)]. The fractional part $n\Phi$ determines the angular position of the n th leaf measured in turn, where Φ denotes the golden ratio. A positive integer n is called a “front number” if $n\Phi < 1/4$ or $n\Phi > 3/4$, otherwise, n is called a “back number”.

The sequence of front numbers (<http://oeis.org/A295085>) is related with some sequences of Kimberling: the sequence of front numbers is the intertwining of A190249 and A190251 and the sequence of back numbers corresponds to A190250.

Burghard Herrmann, Köln, Germany

Visibility in a Pure Model of Golden Spiral Phyllotaxis

As a contribution to our understanding of lattices the talk summarizes the paper “Visibility in a pure model of golden spiral phyllotaxis”. It is published in the current issue of MATHEMATICAL BIOSCIENCES (Share Link: <https://authors.elsevier.com/a/1X9t75pvHBD-A>).

Orli Herscovici, University of Haifa, Haifa, Israel

New Degenerated Bernoulli and Euler Polynomials Arising From Non-Classical Umbral Calculus

We introduce new generalizations of the Bernoulli and Euler polynomials based on the degenerate exponential function and concepts of the Umbral Calculus associated with it. We present generalizations of some familiar identities and connection between these kinds of Bernoulli and Euler polynomials which we have established in our preliminary work.

(Joint work with Toufik Mansour.)

Lin Jiu, Dalhousie University, Halifax, NS, Canada

Bessel Random Walks and Identities for Higher-Order Bernoulli and Euler Polynomials

We consider the study of random walks as a technique to obtain non-elementary identities for higher-order Euler and Bernoulli polynomials. In the case of a one-dimensional linear reflected Brownian motion, considering the successive hitting times of uniformly distributed levels in $[0, 1]$ yields non-trivial expressions for higher-order Euler polynomials. These results are also interpreted as a stochastic sum decomposition due to Klebanov. Analogous results in the case of a 3-dimensional Bessel process yield non-elementary expressions about higher-order Bernoulli polynomials.

(Joint work with Christophe Vignat.)

Bir Kafle, Purdue University Northwest, Westville/Hammond, IN, USA

Pell Numbers of the Form $2^a + 3^b + 5^c$

The Pell sequence $(P_n)_{n \geq 0}$, Pell-Lucas sequence $(Q_n)_{n \geq 0}$ and the associated Pell sequence $(q_n)_{n \geq 0}$ are defined by the same binary recurrences

$$P_{n+1} = 2P_n + P_{n-1}, \quad Q_{n+1} = 2Q_n + Q_{n-1} \quad \text{and} \quad q_{n+1} = 2q_n + q_{n-1},$$
with the initial terms $P_0 = 0, P_1 = 1, Q_0 = Q_1 = 2$ and $q_0 = q_1 = 1$, respectively. The problem of finding Fibonacci, Lucas, or Pell numbers of a particular form has a very rich history. In this talk, we look into P_n, Q_n and q_n as the sum of the three perfect powers of some prescribed distinct bases. In particular, we determine all the solutions of the Diophantine equations

$$P_n = 2^a + 3^b + 5^c, \quad Q_n = 2^a + 3^b + 5^c \quad \text{and} \quad q_n = 2^a + 3^b + 5^c$$

in positive integers (n, a, b, c) , with some restrictions. Our methods involve the linear forms in logarithms of algebraic numbers.

(Joint work with F. Luca and A. Togbé.)

Clark Kimberling, University of Evansville, Evansville, IN, USA
Linear Complementary Equations and Systems

After a brief history of complementary equations, a definition is given for linear complementary equation, with particular attention to examples typified by $a_n = a_{n-1} + a_{n-2} + b_n$, where (b_n) is the complement of (a_n) in the set N of positive integers, and $a_n/a_{n-1} \rightarrow (1 + \sqrt{5})/2$. Also introduced are systems of equations whose solutions are sequences that partition N . An example is the system defined recursively by $a_n =$ least new k , $b_n =$ least new k , and $c_n = a_n + b_n$, where “least new k ”, also known as “mex”, is the least integer in N not yet placed. The sequence (c_n) in this example is the anti-Fibonacci sequence, A075326 in the Online Encyclopedia of Integer Sequences.

(Joint work with Peter J. C. Moses.)

Karyn McLellan, Mount Saint Vincent University, Halifax, NS, Canada
A Problem on Generating Sets Containing Fibonacci Numbers

At the Sixteenth International Conference on Fibonacci Numbers and Their Applications the following problem was posed:

Let S be the set generated by these rules: Let $1 \in S$ and if $x \in S$, then $2x \in S$ and $1 - x \in S$, so that S grows in generations: $G_1 = \{1\}$, $G_2 = \{0, 2\}$, $G_3 = \{-1, 4\}$, \dots

Prove or disprove that each generation contains at least one Fibonacci number or its negative.

We will show that every integer k can be found in some G_i and will disprove the above statement by finding an expression for the generation index i for any given k . We will use a variety of recurrence sequences including the dragon curve sequence, properties of binary numbers, and a computer calculation to find numerous counterexamples.

(Joint work with Danielle Cox)

Steven J. Miller, Williams College, Williamstown, MA, USA
From Monovariants to Zeckendorf Decompositions and Games

Zeckendorf’s Theorem states that every positive integer has a unique decomposition as a sum of non-adjacent Fibonacci numbers; this has been generalized to many other recurrences. We show by looking at appropriate monovariants that these decompositions have the fewest summands possible. We use this perspective to analyze a new two-player game on Fibonacci decompositions, and provide a non-constructive

proof that Player Two always has a winning strategy. As time permits we will discuss generalizations and open problems.

Antara Mukherjee, The Citadel, Charleston, SC, USA

The Geometric Interpretation of Some Fibonacci Identities in the Hosoya Triangle

The Hosoya triangle is a triangular array (like the Pascal triangle) where the entries are products of Fibonacci numbers. The symmetry present in the Hosoya triangle helps us explore several patterns and find new identities. In this talk we give a geometric interpretation -using the Hosoya triangle- of several Fibonacci identities that are well known algebraically. For example, we discuss geometric proofs of Cassini, Catalan, and Johnson identities. We also extend some properties from Pascal triangle to the Hosoya triangle. For instance, we generalize the hockey stick property, the T-stick identities – that were originally given in terms of binomial coefficients – to identities for Fibonacci numbers.

(Joint work with R. Flórez and R. Higueta.)

Kouichi Nakagawa, Saitama University, Saitama, Japan

Exact Periodicity of Generalized Fibonacci and Tribonacci Sequence

Let $\{G_n(a, b)\}$ be the generalized Fibonacci sequence, where $G_0 = a$, $G_1 = b$, and $G_n = G_{n-1} + G_{n-2}$, $n \geq 2$, and let $\{T_n(a, b, c)\}$ be the generalized Tribonacci sequence, where $T_0 = a$, $T_1 = b$, $T_2 = c$ and $T_n = T_{n-1} + T_{n-2} + T_{n-3}$, $n \geq 3$. (Thus $G_n(0, 1)$ is the n th Fibonacci number, $G_n(2, 1)$ is the n th Lucas number, $T_n(0, 0, 1)$ is the n th Tribonacci number (or n th Fibonacci 3-step number) and $T_n(3, 1, 3)$ is the n th Lucas 3-step number).

D. D. Wall showed that the generalized Fibonacci sequence is simply periodic when taken modulo m . (For example, in the case of the original Fibonacci sequence the length of period (mod 10) is 60, and (mod 100) it is 300, and so on.) C. C. Yalavigi showed that the generalized Tribonacci sequence mod m is simply periodic as well. (For example, in the case of the original Tribonacci sequence the length of period (mod 10) is 124 and (mod 100) is 1240, and so on.) However, the simply periodic sequences do not necessarily have the smallest period. (For example, the generalized Tribonacci sequences (mod 10), $(a, b, c) = (3, 1, 3)$, $(1, 7, 9)$, \dots have a period of 31, $(a, b, c) = (0, 1, 0)$, $(3, 2, 1)$, \dots have a period of 62 and $(a, b, c) = (0, 0, 1)$, $(2, 3, 5)$, \dots have a period of 124.) Hence we compute the periods (mod 10^d) up to $d = 4$ for all generalized Fibonacci sequences and up to $d = 3$ for all Tribonacci

sequences by experimental mathematics and analyze the relationships between periodic groups observed.

(Joint work with Rurika Sudo.)

Sam Northshield, SUNY-Plattsburgh, Plattsburgh, NY, USA

Re³ counting the Rationals

In 1999, Neil Calkin and Herbert Wilf wrote their charming “Re-counting the rationals” which gave an explicit bijection between the positive integers and the positive rationals: namely, $n \mapsto a_{n+1}/a_n$ where a_n is defined by $a_1 = 1$, $a_{2n} = a_n$, and $a_{2n+1} = a_{n+1} + a_n$. Alternatively, a_{n+1} is the number of hyperbinary representations of n or, in still another way,

$$a_{n+1} = a_n + a_{n-1} - 2(a_{n-1} \bmod a_n).$$

Expressing this in another way, for $f(x) := 1 + 1/x - 2\{1/x\}$ where $\{x\}$ denotes the fractional part of x , the sequence $1, f(1), f(f(1)), f(f(f(1))), \dots$ is a list of all of the positive rationals.

We will discuss the facts that iterates of $2 + 2/x - 4\{1/x\}$ starting at 2 also cover the positive rationals as do the iterates of $3 + 3/x - 6\{1/x\}$ starting at 3. That is, the iterates of $cf(x)$, starting at c , cover the positive rationals when $c = 1, 2, 3$. Surprisingly, $c = 1, 2, 3$ are the only numbers for which this is true.

I’ll sketch some of the proofs; they involve, among other things, “negative” continued fractions, Chebyshev polynomials, Euler’s totient function, arrangements of circles, and the generalized Stern sequences

$$x_{n+1} = \sqrt{c} \cdot x_n + x_{n-1} - 2(x_{n-1} \bmod (\sqrt{c} \cdot x_n)).$$

I will also discuss some remarkable properties of these latter sequences.

Prapanpong Pongsriiam, Silpakorn University, Faculty of Science, Nakhon Pathom, Thailand

Fibonacci and Lucas Numbers Which Have Exactly Three Prime Factors and Some Unique Properties of F_{18} and L_{18}

Let F_n and L_n be the n th Fibonacci and Lucas numbers, respectively. Let $\omega(n)$ be the number of prime factors of n , $d(n)$ the number of positive divisors of n , $A(n)$ the least positive reduced residue system modulo n , and $\ell(n)$ the length of longest arithmetic progressions contained in $A(n)$. In the occasion of attending the 18th Fibonacci Conference, we will show some results concerning $\omega(F_n)$, $\omega(L_n)$, $d(F_n)$, and $d(L_n)$ which reveal a unique property of F_{18} and L_{18} . We also find the solutions to the equation $\ell(n) = 18$ and show a connection

between them and F_{18} . Some examples and numerical data will also be presented.

J. C. Saunders, University of Waterloo, Waterloo ON, Canada
On (a, b) Pairs in Random Fibonacci Sequences

We examine the random Fibonacci tree, which is an infinite binary tree with non-negative integers at each node. The root consists of the number 1 with a single child, also the number 1. We define the tree recursively in the following way: if x is the parent of y , then y has two children, namely $|x - y|$ and $x + y$. This tree was studied by Benoit Rittaud who proved that any pair of integers a, b that are coprime occur as a parent-child pair infinitely often. We extend his results by determining the probability that a random infinite walk in this tree contains exactly one pair $(1, 1)$, that being at the root of the tree. Also, we give tight upper and lower bounds on the number of occurrences of any specific coprime pair (a, b) at any given fixed depth in the tree.

(Joint work with Kevin Hare.)

Susanna Spektor, Brock University, St. Catharines, ON, Canada
On a ψ_1 -Norm Estimate of Sums of Dependent Random Variables Using Simple Random Walks on Graphs

We obtained a ψ_1 estimate for the sum of Rademacher random variables under condition that they are dependent.

Paul K. Stockmeyer, The College of William & Mary, Williamsburg, VA, USA

Discovering Fibonacci Numbers, Fibonacci Words, and a Fibonacci Fractal in the Tower of Hanoi

The Tower of Hanoi puzzle, with three pegs and n graduated discs, was invented by Edouard Lucas in 1883, writing under the name of Professor N. Claus. A simple question about relative distances between various regular states of this puzzle has led to the discovery of a new occurrence of Fibonacci numbers, a new illustration of the finite Fibonacci words, and a fractal of Hausdorff dimension $\log_2(\phi)$, where ϕ is the golden ratio $\frac{1+\sqrt{5}}{2}$.

(joint work with Andreas M. Hinz, LMU München, Germany.)

Elif Tan, Ankara University, Ankara, Turkey
A Note on Conditional Divisibility Sequences

A sequence of rational integers $\{a_n\}$ is said to be a *divisibility sequence (DS)* if $m \mid n$ whenever $a_m \mid a_n$ and it is said to be a *strong divisibility sequence (SDS)* if $\gcd(a_m, a_n) = a_{\gcd(m,n)}$. These sequences are of particular interest because of their remarkable factorization properties and usage in applications, such as factorization problem, primality testing, etc. The best known examples are Fibonacci sequence, Lucas sequence, Lehmer sequence, Vandermonde sequences, resultant sequences and their divisors, elliptic divisibility sequences, etc.

In this talk, we consider the conditional recurrence sequence $\{q_n\}$ is one in which the recurrence satisfied by q_n depends on the residue of n modulo some integer $r \geq 2$. If the conditional sequence $\{q_n\}$ is also a divisibility or strong divisibility sequence, we call it as a *conditional divisibility* or *conditional strong divisibility sequence*. We investigate and find some families of the conditional divisibility and the conditional strong divisibility sequences.

(Joint work with Murat Şahin.)

Thotsaporn ‘Aek’ Thanatipanonda, Mahidol University International College, Nakornphathom, Thailand
Statistics of Domino Tilings on a Rectangular Board

It is well known the Fibonacci sequence, F_n , is the number of ways to cover a 2-by- $(n-1)$ board using only the horizontal(H) or vertical(V) 2-by-1 dominos. It is natural to generalize this idea to a rectangular m -by- n board where m is a fixed number and n is symbolic. We can try harder and compute the mixed moment $S[V^a H^b]$ for fixed non-negative integers a, b but general m, n . After all these moments are computed, we will gain an information of the distribution of these statistics as well. Note that the Fibonacci numbers and generalization are the cases where $a = b = 0$ i.e. the zero moment.

Christophe Vignat, Université d’Orsay, Orsay, France
Finite Generating Functions for the Sum-of-Digits Sequence

I will show some results about finite generating functions associated with the sequence $\{s_b(n)\}$, where $s_b(n)$ is the sum of the digits of the representation in base b of the integer n . This sequence has been studied, for example, by J.-P. Allouche and J. Shallit – see their book “Automatic Sequences, Theory, Applications, Generalizations”. Thanks to a general identity that relates the sequence $\{s_b(n)\}$ to the

finite difference operator, we obtain, for example, an explicit expression for a Hurwitz-type generating function related to this sequence. Our generalizations also include links to some Lambert series and to infinite products related to the sequence $s_b(n)$.

(Joint work with T. Wakhare.)

Tanay Wakhare, University of Maryland, College Park, MD, USA
Structural Identities for Multiple Zeta Values

medskip We revisit some results by Borwein et al. about Multiple Zeta Values and show that they can be extended to an arbitrary, possibly finite, sequence of numbers. Specializing these sequences as the zeros of special functions gives us some new results about Bessel and Airy Multiple Zeta values. In the Bessel case, specializing the argument to $\nu = 1/2$ allows us to recover the classical results by Borwein.

(Joint work with C. Vignat.)

William Webb, Washington State University, Pullman, WA, USA
What Makes A “Nice” Identity?

There are probably thousands of known identities involving recurrence sequences. We will suggest one way to judge whether an identity is particularly simple, or nice, or maybe unexpected. We begin with a reminder that often the easiest way to prove many of the basic properties of recurrence sequences, including proving identities, is viewing recurrences as elements of vector spaces. By looking at the dimensions of these vector spaces we can show why some identities are rather ordinary and others much more unexpected. We end by showing how these techniques can be used to prove a conjecture by R. S. Melham.

William Webb, Washington State University, Pullman, WA, USA
Proving Identities in Arbitrary Fields

Most of the known identities involve the Fibonacci numbers or possibly other recurrence sequences in the ring of integers. However, the natural place to study them is in the complex field, since we often need to express a recurrence as a generalized power sum in terms of powers of the roots of the associated characteristic polynomial. Since we need roots of polynomials it is easier to work over an algebraically closed field. It is often the case that an identity (or other result) is first proved for the Fibonacci numbers, then maybe for the Lucas numbers, Pell numbers, other second order integer sequences, Fibonacci polynomials etc. However, it is possible to prove all of these cases, as well

as for recurrence sequences in much more general algebraically closed fields, all at once. Since the results are much more general, the proofs may be more tedious. Many proof techniques can be used, but some, such as combinatorial proofs which usually involve quantities counted by integers, may not be applicable. We give several examples of such general identities, such as:

If a second order recurrence sequence satisfies the recurrence $u_{n+2} = au_{n+1} + bu_n$, then

$$\sum_{j=0}^n u_j = \frac{bu_n + u_{n+1} - 1}{a + b - 1}.$$

This is of course not a new identity, but we note that it is true regardless of whether the parameters a and b and hence the recurrence sequence itself are integers, polynomials, power series, p -adic numbers etc.

(Joint work with Nathan Hamlin.)

Paul Young, College of Charleston, Charleston, SC, USA

Global Series for Zeta Functions

We give two general classes of everywhere-convergent series for Barnes generalization of Hurwitz zeta functions, which involve Bernoulli polynomials of the second kind and weighted Stirling numbers. These series are also valid p -adically, and yield several identities and series for zeta values and Stieltjes constants which are valid in both real and p -adic senses.

Paul Young, College of Charleston, Charleston, SC, USA

The Power of 2 Dividing a Generalized Fibonacci Number

Let T_n denote the generalized Fibonacci number of order k defined by the recurrence $T_n = T_{n-1} + T_{n-2} + \cdots + T_{n-k}$ for $n \geq k$, with initial conditions $T_0 = 0$ and $T_i = 1$ for $1 \leq i < k$. Motivated by some recent conjectures of Lengyel and Marques, we establish the 2-adic valuation of T_n , settling one conjecture affirmatively and one negatively. We discuss the computational issues that arise and applications to Diophantine equations involving (T_n) .

Osman Yürekli, Ithaca College, Ithaca, NY, USA

A Pascal-Like Triangle From a Special Function

This presentation is devoted to a new Pascal-like triangle appearing unexpectedly from a sequence of polynomials obtained from the

derivatives of the special function Dawson's integral which is defined by the integral

$$\text{daw}(x) = \int_0^x \exp(y^2 - x^2) dy.$$

We investigate the properties of the Pascal-like triangle and its applications. In addition, we discuss a generalization for the triangle and Dawson's integral. It is also possible to obtain new Fibonacci-like sequences from the triangle and its generalizations.

The Paul Bruckman Prizes

The Fibonacci Association is pleased to announce the establishment of the Paul Bruckman Prize program pursuant to which prizes of \$1000 will be awarded for papers which develop a new approach or expand results in the area of generalized Fibonacci numbers and related areas of mathematics. Guidelines for topics which would qualify for submission can be obtained by email from: fibonacciassociation@gmail.com.

Two prizes will be awarded in each even-numbered year commencing with 2016. One prize will be awarded to a paper appearing in the last two years in the Fibonacci Quarterly. The second prize will be awarded to a paper presented at the Fibonacci Association biannual conference in the relevant year.

The prize is named in honor of Paul Bruckman who had a long and distinguished association with Fibonacci numbers as described in more detail in A Tribute to Paul S. Bruckman published in *The Fibonacci Quarterly*, **49.3** (2011), 281. The program is funded by a grant from George A. Hisert of Berkeley, California. George started his career as a mathematician, but for various reasons he then chose to become a lawyer. Upon his retirement from practicing law, he returned to mathematics with a particular interest in Fibonacci numbers. That interest was furthered through several interactions with Paul Bruckman. Three papers, authored by George, were published in the *JP Journal of Algebra, Number Theory and Applications*, Volumes 24, 28, and 31.

The two prizes will be awarded at the Eighteenth International Conference on Fibonacci Numbers and their Applications scheduled to be held in Halifax, Nova Scotia, Canada, in the summer of 2018. Papers will be judged by a panel chosen from the Fibonacci Association's Board of Directors or its Editorial Board. In the event that there is no paper submitted which the panel considers as qualifying for a prize, no prize will be given in that category for that year.

The aim of the prize is to recognize researchers who are early in their careers. Special consideration will be given to papers where at least one of the co-authors is 40 or younger. Age will be determined as of the date a paper is submitted.

The winners of the 2018 George Bruckman Prizes will be announced during the conference dinner on Thursday, July 5, 2018.

The Fibonacci Portrait

No true portrait of Leonardo of Pisa (Fibonacci) seems to be known, but probably the most impressive one of the very few imagined portraits of Fibonacci is the photograph that was chosen for the cover of this program and also for the commemorative mugs. This photograph was taken by the late Frank Johnson (1934–2016), a regular attendee of the International Fibonacci Conferences, well-known in the Fibonacci community, and a friend to many.

This portrait has been reproduced with the kind permission of Frank’s widow, Marjorie Bicknell-Johnson. Leading up to the current conference, Marjorie wrote the following about the history of this photograph:

This portrait was taken by Frank Johnson in 1978 as a gift for Dr. Verner E. Hoggatt, Jr., co-founder of the Fibonacci Quarterly. The statue, larger than life, stood on a pedestal, and its head was at least fifteen feet above the ground. Frank climbed onto rickety scaffolding and leaned out to capture this picture; I was afraid he’d fall and break his neck. Frank accompanied me at all but two Fibonacci Conferences.



Meanwhile the statue has been cleaned up and has been moved to a sheltered location. It will therefore have lost its greenish tinge which, to me, makes this portrait so appealing. More about the statue, its history, and its current location can be found in Keith Devlin’s second book about Fibonacci, namely “Finding Fibonacci: The Quest to Rediscover the Forgotten Mathematical Genius Who Changed the World”, Princeton University Press, 2017.

Choices for Lunch

Monday, July 2:

This being a holiday, almost everything on campus will be closed, with the exception of the **Howe Dining Hall**; see below. Since Monday's lunch break is 2 hours long (as opposed to 1 1/2 hours Tuesday–Friday), some of the closer off-campus places may also be an option; see the following page.

Tuesday–Friday:

We can recommend the following close-by lunch places on campus:

1. The Pub in the University Club, downstairs (see the campus map – it's across the “Studley Quad”).

Very popular among faculty and staff (including your organizers). Fast service, decent food, and an “overflow room” if it gets too busy. A variety of sandwich choices as well as soups and hot dishes, including vegetarian. Typically \$10–\$15.

2. The Dining Room in the University Club, main floor.

Same food choices as in the Pub, but in a more “genteel” dining room setting. It will take more time, but the 90 minute breaks will be sufficient.

3. The Howe Dining Hall, located in the Howe Hall Residence (see campus map – the purple “2” on Coburg Road.)

Open for lunch, 11 am – 2 pm. \$10 + tax. One of the main dining halls to serve students in residence, but also off-campus students, as well as visitors during the Summer months. Other meals served there daily: Breakfast: 7–10am – \$8.50; Dinner: 5–7:30pm – \$14.00.

4. Student Union Building; certainly the closest, right across from the conference venue.

A choice of different fast food places, including the iconic Canadian “Tim Horton's”. Lots of space.

5. Coburg Coffee House, just off campus, at the corner of Coburg Road and Henry Street.

Open 7:00am–7:00pm; some days longer. Very popular among students and faculty. Offers simple breakfasts, sandwiches and baked goods. May get busy at lunch, but the 90 minute breaks will suffice.

Off-Campus Restaurants

There is a small cluster of restaurants not too far from Campus, near Robie Street, where Coburg Road changes into Spring Garden Road. It's approximately 7 minutes to walk from the conference venue (the McCain Building).

Efes Turkish Cuisine, 5986 Spring Garden Road

A popular Turkish restaurant, considered quite good.

Mary's Place Cafe II, 5982 Spring Garden Road

"Old-school, unfussy cafe offering all-day breakfast and Canadian/Middle Eastern lunch plates".

On the same block: A few more fast-food, Chinese and Pizza places.

Mappatura Bistro, 5883 Spring Garden Road

A few minutes further down the street, on the other side. Relatively new, and we haven't been there yet. See mappaturabistro.ca.

Downtown and Waterfront Restaurants

Too many and too varied to make particular recommendations. Please see a tourist guide. However, we can recommend three restaurants to which we routinely take visitors:

Curry Village, 1569 Dresden Row (in the downtown core)

"Comfortable, long-running nook providing traditional Indian meals, lunch specials and outdoor seating." <http://www.curryvillage.ca>

Cha Baa Thai Restaurant, 1546 Queen Street (downtown)

"Fresh Ingredients and Authentic Taste". <http://chabaathairestaurant.ca>

Salty's Seafood Restaurant, 1877 Upper Water Street

Located in the "Historic Properties", right on the waterfront. For a table in the upstairs dining room, which offers a great view of the harbour, a reservation is recommended. In addition to traditional seafood, a variety of non-seafood items are also on the menu. <http://www.saltys.ca>.

Books on Display

There will be a small display of books by authors who are past or current conference participants, and some other relevant books. Most will be display copies. Please have a look at them and consider ordering or purchasing some of them.

Arthur Benjamin and Jennifer Quinn:

Proofs that Really Count: The Art of Combinatorial Proof
MAA Press, 2003.

Arthur Benjamin:

The Magic of Math: Solving for x and Figuring Out Why
Basic Books, 2016.

Arthur Benjamin and Michael Shermer:

Secrets of Mental Math: The Mathemagician's Guide to Lightning Calculation and Amazing Math
Three Rivers Press, 2006.

Marc Chamberland:

Single Digits: In Praise of Small Numbers
Princeton University Press, 2017.

Keith Devlin:

Finding Fibonacci: The Quest to Rediscover the Forgotten Mathematical Genius Who Changed the World
Princeton University Press, 2017.

Richard Dunlap:

The Golden Ratio and Fibonacci Numbers
World Scientific, 1997.

Michael J. Jacobson, Jr. and Hugh C. Williams:

Solving the Pell Equation
CMS Books in Mathematics, Springer, 2009.

George M. Phillips: *Two Millennia of Mathematics. From Archimedes to Gauss*

CMS Books in Mathematics, Springer, 2000.

George M. Phillips: *Interpolation and Approximation by Polynomials*

CMS Books in Mathematics, Springer, 2003.

Anthony G. Shannon and Jean V. Leyendekkers

The Fibonacci Numbers and Integer Structure
Nova Science Publ., 2018.

Bookshops in Halifax

There is only one decent place for new books in central Halifax:

Bookmark, 5686 Spring Garden Road

Located across the street from the Lord Nelson Hotel, and very close to the Public Gardens, this is a small but well-stocked bookshop. <https://bookmarkreads.ca/>

Only a few blocks further, at the corner of Spring Garden Road and Queen Street, you will find a new Halifax landmark, the Public Library. Well worth a visit for its architecture, contents, and two coffee shops, one of them on the top floor which, by the way, offers an excellent view.

Second-hand Bookshops

For those who like second-hand and antiquarian books, we can recommend the following shops. Their owners and staff are all knowledgeable and passionate about books.

Schooner Books, 5378 Inglis Street

A Halifax institution for over 40 years, located in the South End, not far from the Port of Halifax and also within walking distance of Point Pleasant Park. Well worth a visit. <http://www.schoonerbooks.com>

Dust Jacket Books And Treasures, 1505 Barrington St,

A hidden treasure, located in the basement of the “Maritime Centre” highrise building, right downtown, where Spring Garden Road meets Barrington Street, very close to St. Mary’s Basilica and the Old Burying Ground.

The Last Word Bookstore, 2160 Windsor St.

This is the smallest of the second-hand bookstores, but still has plenty to offer and is worth a visit. It’s only a few minutes by foot from the Atlantica Hotel.

John W. Doull, Bookseller, 122 Main St., Dartmouth

To visit this amazing bookshop you would need a car, or time and patience with a fairly long bus ride. Don’t be fooled by the drab exterior and ugly (though safe) neighbourhood. This shop has a huge selection of books, and very knowledgeable staff. A book lover’s paradise. <http://www.doullbooks.com>

Second-hand Mathematics Books

Last, but not least, in the Mathematics & Statistics Department (Chase Building) there is a huge collection of second-hand mathematics books, with some statistics, computer science, and some other sciences. This is a fundraising initiative. See

<https://www.mathstat.dal.ca/~dilcher/oldbooks.html>

and talk with Karl Dilcher if you're interested in any of the books listed. The books are not publicly accessible, but I'll try to find a time during the conference to provide access for conference participants.